

Quasi-Partial sums of the generalized Bernard integral of certain analytic functions

K. O. Babalola

Department of Mathematics, University of Ilorin, Ilorin, Nigeria
 e-mail: ummusalamah.kob@unilorin.edu.ng

Abstract

In this short note we extend a result of Jahangiri and Farahmand [5] concerning functions of bounded turning to a more general class of functions

1.0 Introduction

Let C be the complex plane. Denote by A the class of functions:

$$f(z) = z + a_2 z^2 + \dots \tag{1.1}$$

which are analytic in the unit disk $E = \{z : |z| < 1\}$

In [5] Jahangiri and Farahmand studied the partial sums of the Libera integral of the class $B(\beta)$, which consist of functions in A satisfying $\text{Re } f'(z) > \beta, 0 \leq \beta < 1$. Functions in $B(\beta)$ are called functions of bounded turning. It is known that functions of bounded turning are generally univalent and close-to-convex in the unit disk. In particular they proved that the m th partial sums

$$F_m(z) = z + \sum_{k=2}^m \frac{2}{k+1} a_k z^k \tag{1.2}$$

of the Libera integral

$$F(z) = \int_0^z f(t) dt \tag{1.3}$$

is also of bounded turning. Their result was stated as:

Theorem A.

$$\text{If } \frac{1}{4} \leq \beta < 1 \text{ and } f \in B(\beta), \text{ then } F_m \in B\left(\frac{4\beta-1}{3}\right)$$

Earlier and Owa [6] have proved that if $f \in A$ is univalent in E , then the partial sum $F_m(z)$ is starlike in the subdisk $|z| < \frac{3}{8}$, the number $\frac{3}{8}$ being the best possible.

The result of Jahangiri and Farahmand [5] naturally leads to inquisition about a more general class of functions (including $B(\beta)$ as a special case), which was introduced in [7] by Opoola, has been studied extensively in [2]. This is the class $T_n^a(\beta)$ consisting of functions $f \in A$ which satisfy the inequality;

$$\text{Re } \frac{D^n f(z)^a}{\alpha^n z^a} > \beta \tag{1.4}$$

where $\alpha > 0$ is real, $0 \leq \beta < 1$, D^n ($n \in N_0 = \{0, 1, 2, \dots\}$) is the Salagean derivative operator defined as

$$D^n f(z) = D[D^{n-1} f(z)] = z[D^{n-1} f(z)]' \tag{1.5}$$

with $D^0 f(z) = f(z)$ and powers in (1.4) meaning principal values only. Obverse that the geometric condition (1.4) slightly modifies the only given originally in [7] (see [2]). Onverse also that the class $B(\beta)$ corresponds to $n = \alpha = 1$.

In a recent work we considered the generalized Bernardi integral operator given by

$$F(z)^\alpha = \frac{\alpha + c}{z^c} \int_0^z t^{c-1} f(t)^\alpha dt, \quad \alpha + c > 0 \tag{1.6}$$

and sharpened and extended many earlier results concerning closure, under the integral, of several classes of functions. In the present paper we define a concept of quasi-partial sums and follow a method of Jahagiri an Farahmand [5] to extend their result (Theorem A) to the class $T_n^\alpha(\beta)$.

As we noted in [1], the binomial expansion of (1,1) gives

$$f(z)^\alpha = z^\alpha + \sum_{k=2} a_k(\alpha) z^{\alpha+k-1} \tag{1.7}$$

where $a_k(\alpha)$ is a polynomial depending on the coefficients of $f(z)$ and the index α . Hence

$$F(z)^\alpha = z^\alpha + \sum_{k=2} \frac{\alpha + c}{\alpha + c + k - 1} a_k(\alpha) z^{\alpha+k-1} \tag{1.8}$$

and we define the m th quasi-partial sums of the integral (1.6) as follows

$$F_m(z)^\alpha = z^\alpha + \sum_{k=2}^m \frac{\alpha + c}{\alpha + c + k - 1} a_k(\alpha) z^{\alpha+k-1} \tag{1.9}$$

In the next section we state the preliminary results.

2.0 Preliminary Results

We will require the following lemmas.

Lemma 2.1 [3]

Let 0 be a real number and l a positive integer. If $-1 < \gamma \leq A$, then

$$\frac{1}{1+\gamma} + \sum_{k=1}^l \frac{\text{Cos}k\theta}{k+\gamma} \geq 0$$

The constant $A = 4.5678018, \dots$ is the best possible.

Lemma 2.2

For $z \in E, -1 < \gamma \leq A = 4.5678018, \dots$, $\text{Re} \left(\sum_{k=1}^l \frac{z^k}{k+\gamma} \right) \geq -\frac{1}{1+\gamma}$

Proof

Let $z = re^{i\theta}$ where $0 \leq r < 1, 0 < |\theta| \leq \pi$. Then by De Moivre's law and the minimum principle for harmonic functions $\text{Re} \left(\sum_{k=1}^l \frac{z^k}{k+\gamma} \right) = \sum_{k=1}^l \frac{r^k \text{Cos}k\theta}{k+\gamma} > \sum_{k=1}^l \frac{\text{Cos}k\theta}{k+\gamma}$.

Hence by Abel's Lemma [8, pg 6] and Lemma 2.1 above the conclusion follows. Let P denote

the class of analytic functions of the form

$$p(z) = 1 + c_1z + \dots \tag{2.1}$$

normalized by $p(0) = 1$ and satisfy $\text{Re} p(z) > 0$ in E . The next lemma concerns convolution of analytic functions with functions in P . The convolution (or Hadamard product) of two power series $f(z) = \sum_{k=0}^{\infty} a_k b_k z^k$ and $g(z) = \sum_{k=0}^{\infty} b_k z^k$ (written as $f * g$) is defined as $(f * g)(z) = \sum_{k=0}^{\infty} a_k b_k z^k$.

Lemma 2.3 [4]

Let $p(z)$ be analytic in E and satisfy $p(0) = 1$ and $\text{Re} p(z) > \frac{1}{2}$ in E . For analytic function $a(z)$ in E , the convolution $p * a$ takes values in the convex hull of the image of E under $a(z)$.

3.0 Main Results

Theorem 3.1

Let $f(z)$ given by (1.1) be in the class $T_n^\alpha(\beta)$. Then

$$\text{Re} \frac{D^n F_m(z)^\alpha}{\alpha^n z^\alpha} > 1 - \frac{2(1-\beta)(\alpha+c)}{(\alpha+c+1)}, \quad \alpha+c \leq 4.5678018\dots \tag{3.1}$$

Furthermore if $\beta \geq \frac{1}{2} \frac{(\alpha+c-1)}{(\alpha+c)}$, then $F_m(z)$ belongs to some subclasses of the class $T_n^\alpha(\beta)$

Proof

From (1.7) and the condition (1.4) we have

$$\text{Re} \left\{ 1 + \frac{1}{2(1-\beta)} \sum_{k=2}^{\infty} \left(\frac{\alpha+k-1}{\alpha} \right)^n a_k(\alpha) z^{k-1} \right\} > \frac{1}{2} \tag{3.2}$$

Also from (1.9) we have

$$\frac{D^n F_m(z)^\alpha}{\alpha^n z^\alpha} = 1 + \sum_{k=2}^{\infty} \left(\frac{\alpha+k-1}{\alpha} \right)^n \frac{\alpha+c}{\alpha+c+k-1} a_k(\alpha) z^{k-1} = p(z) * q(z) \tag{3.3}$$

where

$$p(z) = 1 + \frac{1}{2(1-\beta)} \sum_{k=2}^{\infty} \left(\frac{\alpha+k-1}{\alpha} \right)^n a_k(\alpha) z^{k-1}, \tag{3.4}$$

$$q(z) = 1 + 2(1-\beta) \sum_{k=2}^m \frac{\alpha+c}{\alpha+c+k-1} z^{k-1} \tag{3.5}$$

Thus by Lemma 2.3 and the condition (3.1) the geometric quantities $D^n F_m(z)^\alpha / \alpha^n z^\alpha$ takes values in the convex hull of $q(E)$. But

$$\text{Re} q(z) = 1 + 2(1-\beta)(\alpha+c) \text{Re} \left(\sum_{k=1}^{m-1} \frac{z^k}{\alpha+c+k} \right) \tag{3.6}$$

We know from (1.6) that $\alpha + c > 0$. Now suppose $\alpha + c \leq 4.5678018\dots$, then by taking $l = m - 1$ in Lemma 22, the real part of the series on right of (3.6) is greater than $-(\alpha + c + 1)^{-1}$ so that

$$\operatorname{Re} \frac{D^n F_m(z)^\alpha}{\alpha^n z^\alpha} = \operatorname{Re} q(z) > 1 - \frac{2(1-\beta)(\alpha+c)}{(\alpha+c+1)} \quad (3.7)$$

Now observe that the real number $1 - \frac{2(1-\beta)(\alpha+c)}{(\alpha+c+1)}$ is nonnegative only for $\beta \geq \frac{1}{2} \frac{(\alpha+c-1)}{(\alpha+c)}$. Thus only in this case it is clear $F_m(z)$ belongs to some subclasses of the class $T_n^a(\beta)$. This completes the proof. ■

Remark

For $\alpha = 1, c = 0$, the partial sums

$$F_m(z) = z + \sum_{k=2}^m \frac{a_k}{k} z^k \quad (3.8)$$

of the integral

$$F(z) = \int_0^z t^{-1} f(t) dt \quad (3.9)$$

for each $f \in B_n(1)$, belongs to the class $B_n(1)$ in general. More particularly, the partial sum (3.8) of the integral (3.9) of a function of bounded turning in the unit disk is also a function of bounded turning in the unit disk.

4.0 Conclusion

In this paper we defined a new concept of quasi-partial sums of the generalized Bernard integral. We used the new concept to extend an earlier result of Jahangiri and Farahmand [5] concerning functions of bounded turning to a more general class of function.

References

- [1] K. O. Babalola, On the generalized Bernard integral of certain analytic functions defined by the Salagean derivative. *Submitted*.
- [2] K. O. Babalola and T. O. Opoola, Iterated integral transforms of Caratheodory functions and their applications to analytic and univalent functions. *Tamkang J. Math* **37** (4) (Winter 2006), 355 – 366.
- [3] G. Gasper, Nonnegative sums of cosines, ultraspherical and Jacobi polynomials. *J. Math. Anal. Appl* **26** (1969), 60 – 68.
- [4] A. W. Goodman, *Univalent functions*. Vol. 1, Mariner Pub. Co., Tampa, FFL., 1983.
- [5] J. M. jahangiri and K. Farahmand, Partial sums of functions of bounded turning, *J. Ineq. Pure and Appl. Maths.* **4**(4) Art. 79 (2003), 1- 9.
- [6] J.L. Li and S. Owan, On partial sums of Libera integral operator, *J. math. Anal. Appl.* **213** (1997) 444 – 434.
- [7] T. O. Opoola, On a new subcasss of univalent function. *Mathematica (Cluj)* (36) 59 No. 2 (1994), 195 – 200.
- [8] E.C. Titchmarsh, *Theory of functions*. 2nd Edition, Oxford University Press, 1976.